

About recurrence time for a semi-Lagrangian discontinuous Galerkin Vlasov solver

Michel Mehrenberger

IRMA, Université de Strasbourg & projet TONUS (INRIA), France

CIRM, 31.10.2017

Joint work with Laurent Navoret (IRMA)

Introduction : results from Cheng-Knorr, 1976



Recurrence time for
SLDG

M. Mehrenberger

Recurrence time $T_R = \frac{2\pi}{k\Delta v}$ is analyzed for free streaming case...

$$\partial_t f + v \partial_x f = 0, \rho = \int f dv, f(t=0) = A \cos(kx) f_0(v), f_0(v) = \frac{1}{\sqrt{2\pi}} e^{-v^2/2}$$

$$\rho(t, x) = A \Delta v \sum_{j=-M}^{M-1} f_0(v_j) \cos(k(x - (j + 1/2)\Delta v t)) \text{ is } T_R \text{ periodic in time}$$



Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo


Conclusion and
perspectives

Introduction : results from Cheng-Knorr 1976

...and observed for the simulation of the linear Landau damping

$$\partial_t f + v \partial_x f + E \partial_v f = 0, \quad \partial_x E = \rho - 1$$

$$f(t=0) = (1 + A \cos(kx)) f_0(v)$$



Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

- ▶ Numerical artefact for several grid-based solvers
- ▶ How to suppress/reduce the phenomenon ?
 - ▶ randomness of the grid
 - ▶ filtering
 - ▶ PIC method
 - ▶ finer grid
- ▶ Recent works : Einkemmer-Ostermann (2014), Cai-Wang (2017)
- ▶ Eliasson (2001, 2010)
- ▶ Objective here : study the behavior of the recurrence for a *Semi-Lagrangian Discontinuous Galerkin* (SLDG) method

The SLDG method for Vlasov-Poisson : history

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

- ▶ Rossmanith-Seal → Qiu-Shu → Crouseilles-M-Vecil (2010-2011)
- ▶ already ideas of SLDG in Mangeney-Califano-et-al (2002)
- ▶ variant+analysis : Einkemmer-Ostermann (2014)
- ▶ adaptive : Besse-Deriaz-Madaule (2016)
- ▶ no splitting : Cai-Guo-Qiu (2017)
- ▶ other contexts :
 - ▶ general approach : Restelli-Bonaventuro-Sacco (2006)
 - ▶ Guo-Nair-Qiu (2014)
 - ▶ analysis : Bokanowski-Simarmata, Bokanowski-Cheng-Shu (2016)
- ▶ some ref. for DG for Vlasov :
 - ▶ Heath-Gamba-et-al (2012)
 - ▶ Cheng-Christlieb-Zhong, Madaule-Restelli-Sonnendrücker (2014)
 - ▶ analysis : Ayuso-Carillo-Shu (2012)

The SLDG method for Vlasov-Poisson : the scheme

Recurrence time for
SLDG

M. Mehrenberger

► In 1D for free streaming

- polynomial representation on each cell
- exact transport (constant velocity)
- projection on each cell : piecewise polynomial \rightarrow polynomial
- can be interpreted as advection of Gauss points¹

$$x_{j,0}, \dots, x_{j,d} \in]x_{\min} + j\Delta x, x_{\min} + (j+1)\Delta x[$$

and written as

$$\begin{pmatrix} f(t + \Delta t, x_{j,0}) \\ \dots \\ f(t + \Delta t, x_{j,d}) \end{pmatrix} = A(\alpha) \begin{pmatrix} f(t, x_{j,0}) \\ \dots \\ f(t, x_{j,d}) \end{pmatrix} + B(\alpha) \begin{pmatrix} f(t, x_{j+1,0}) \\ \dots \\ f(t, x_{j+1,d}) \end{pmatrix}$$

with $x_{j,\cdot} - v\Delta t = x_{i,\cdot} + \alpha\Delta x$ and $A(\alpha), B(\alpha) \in \mathbb{R}^{d+1} \times \mathbb{R}^{d+1}$

► In 1D \times 1D phase-space for Vlasov-Poisson

- polynomial represented by Gauss points on each rectangular cell
- splitting : succession of
 - 1D advections of Gauss points in x direction (each 1D advection updates its horizontal line)
 - Poisson
 - 1D advections of Gauss points in v direction (each 1D advection updates its vertical line)

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

1. we can use other points similarly

Free streaming for Gauss type mesh

Recurrence time for
SLDG

M. Mehrenberger

The mesh in velocity is given by

$$v_{j,\ell} = (j + \bar{v}_\ell) \Delta v, \ell = 0, \dots, d$$

with $0 < \bar{v}_0 < \dots < \bar{v}_d < 1$ and we solve

$$\partial_t f + v_{j,\ell} \partial_x f = 0$$

► Framework not only for SLDG : what counts here is the mesh

► Do we have a recurrence time ?

► is it formula $T_R = 2\pi/(k\Delta v)$?

- OK for $d = 0$ and $\bar{v}_0 = 1/2$.
- but looking at uniform cases :

- $d = 1$ and $\bar{v}_0 = 1/4, \bar{v}_1 = 3/4, \rightarrow T_R = 2\pi/(k\Delta v/2)$

- $d = 2$ and $\bar{v}_0 = 1/6, \bar{v}_1 = 3/6, \bar{v}_2 = 5/6, \rightarrow T_R = 2\pi/(k\Delta v/3)$

- $\dots \bar{v}_k = (2\ell + 1)/(2d + 2), \ell = 0, \dots, d$

we can hope formula $T_R = 2\pi(d + 1)/(k\Delta v)$

► We will see that formula is between :

$$T_R \simeq 2\pi(\lfloor d/2 \rfloor + 1)/(k\Delta v)$$

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

We get

$$\rho(t, x) = A\Delta v \sum_{j=-M}^{M-1} \sum_{\ell=0}^d \omega_{\ell} f_0(v_{j,\ell}) \cos(k(x - (j + \bar{v}_{\ell})\Delta vt))$$

With symmetry of discrete velocity distribution, we obtain

$$\rho(t, x) = A \cos(kx) h(t),$$

with

$$h(t) = \Delta v \sum_{j=-M}^{M-1} \sum_{\ell=0}^d \omega_{\ell} f_0(v_{j,\ell}) \cos(k(j + \bar{v}_{\ell})\Delta vt)$$

We look the expression for $t = t_m = \frac{2\pi m}{k\Delta v}$, $m \in \mathbb{N}^*$

$$h(t_m) = \sum_{\ell=0}^d \omega_{\ell} \cos(2\pi m \bar{v}_{\ell}) \Delta v \sum_{j=-M}^{M-1} f_0(v_{j,\ell})$$

[Introduction](#)[Cheng-Knorr, 1976](#)[Comments and objective](#)[The SLDG method for
Vlasov-Poisson](#)[history](#)[The scheme](#)[Free streaming for
Gauss type mesh](#)[Recurrence & quadrature](#)[Trigonometric quadrature](#)[Numerical results](#)[free streaming](#)[linear Landau damping](#)[Plasma echo](#)[Conclusion and
perspectives](#)

We look for a quadrature rule such that

$$\sum_{\ell=0}^d \omega_{\ell} \cos(2\pi m \bar{v}_{\ell}) = \int_0^1 \cos(2\pi m v) dv,$$

for all $m \leq n$ with n as high as possible.

- Possible for $n = \lfloor d/2 \rfloor$ and

$$\omega_{\ell} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \prod_{j=0, j \neq \ell}^{\lfloor d/2 \rfloor} \frac{\cos(v) - \cos(2\pi \bar{v}_j)}{\cos(2\pi \bar{v}_{\ell}) - \cos(2\pi \bar{v}_j)} dv, \quad \ell = 0, \dots, d.$$

- Optimal points are uniform and lead to $n = d$ (like Gauss integration)
- for Gauss points, weights remain positive until $d = 18$.
- recent references : Peherstorfer, 2011 ; Austin, PhD 2016

⇒ Instead of using classical *polynomial* integration quadrature rule,
we can use this *trigonometric* integration

[Introduction](#)[Cheng-Knorr, 1976](#)[Comments and objective](#)[The SLDG method for
Vlasov-Poisson](#)[history](#)[The scheme](#)[Free streaming for
Gauss type mesh](#)[Recurrence & quadrature](#)[Trigonometric quadrature](#)[Numerical results](#)[free streaming](#)[linear Landau damping](#)[Plasma echo](#)[Conclusion and
perspectives](#)

Numerical results for free streaming

$d = 2$ (gauss points)

	classical	new
h_1	0.02245	0
h_2	0.52996	0.51916
h_3	0.73652	0.73047

$d = 3$ (gauss points)

	classical	new
h_1	$-1.0678 \cdot 10^{-3}$	0
h_2	-0.12573	-0.12664
h_3	-0.74097	-0.74155
h_4	-0.33811	-0.33830

$d = 4$ (gauss points)

	classical	new
h_1	$3.0803 \cdot 10^{-5}$	0
h_2	0.016669	0
h_3	0.30402	0.30657
h_4	0.79870	0.80082
h_5	-0.010458	-0.018835

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

$d = 5$ (gauss points)

	classical	new
h_1	$-5.9704 \cdot 10^{-7}$	0
h_2	$-1.4223 \cdot 10^{-3}$	0
h_3	-0.071038	-0.073117
h_4	-0.50843	-0.50769
h_5	-0.67454	-0.67455
h_6	0.30953	0.30852

$d = 6$ (gauss points)

	classical	new
h_1	$8.3166 \cdot 10^{-9}$	0
h_2	$8.5037 \cdot 10^{-5}$	0
h_3	0.010947	0
h_4	0.17907	0.18741
h_5	0.66873	0.66483
h_6	0.40512	0.40575
h_7	-0.41364	-0.41456

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

- ▶ Gauss quadrature (classical) already good
- ▶ Gauss quadrature becomes better when d increases
- ▶ small improvement for new trigonometric quadrature
- ▶ do not use when $d > 18$ (negative weights)
- ▶ T_R coherent with result of Madaule-Restelli-Sonnendrücker (2014)

Gauss-Lobatto points

The analysis can be extended to Gauss-Lobatto points

Example : $d = 2$, the mesh is uniform :

$$x_0 = 0, x_1 = 1/2, x_2 = 1$$

- ▶ Trigonometric interpolation corresponds to trapezoidal formula :

$$\omega_0 = \omega_2 = 1/4, \omega_1 = 1/2$$

- ▶ Gauss-Lobatto quadrature is Simpson rule :

$$\omega_0 = \omega_2 = 1/6, \omega_1 = 2/3$$

$d = 2$ (gauss-lobatto points)

	classical	new
h_1	$1/3$	0
h_2	1	1
h_3	$1/3$	0
h_4	1	1

Numerical results for free streaming

$d = 4$ (gauss-lobatto points)

	classical	new
h_1	$1.3174 \cdot 10^{-3}$	0
h_2	0.14855	0
h_3	0.79651	0.93393
h_4	0.25734	0.12776
h_5	-0.10028	-0.11932

$d = 6$ (gauss-lobatto points)

	classical	new
h_1	$6.9354 \cdot 10^{-7}$	0
h_2	$1.6300 \cdot 10^{-3}$	0
h_3	0.079357	0
h_4	0.542841	0.663454
h_5	0.64533	0.51383
h_6	-0.34435	-0.22468
h_7	0.15274	$2.1373 \cdot 10^{-3}$


positive weights until $d = 7$.

Numerical results for linear Landau damping

Recurrence time for
SLDG

M. Mehrenberger

$$N = 16, v_{\max} = 4, A = 10^{-2}, k = 0.5, 2\pi/(k\Delta v) \simeq 25.13$$



Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for linear Landau damping

$N = 64$, $v_{\max} = 8$, $A = 10^{-6}$, $k = 0.5$, $2\pi/(k\Delta v) \simeq 50.26$

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for Vlasov-Poisson

history

The scheme

Free streaming for Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature



Numerical results

free streaming

linear Landau damping



Plasma echo

Conclusion and perspectives



Numerical results for plasma echo

$N = 600$, $v_{\max} = 8$, $A = 10^{-3}$, $k_1 = \frac{12\pi}{100}$ perturbation with $k_2 = \frac{24\pi}{100}$ at time $t = 200$ is added. $2\pi/(k_1 \Delta v) = 625$



Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

- ▶ Conclusion
 - ▶ link between recurrence and quadrature
 - ▶ introduction of trigonometric quadrature for non uniform velocity discretization
 - ▶ analysis for free streaming
 - ▶ numerical results for Landau damping
- ▶ Perspectives/possible applications
 - ▶ velocity semi-discretization method (N. Pham, PhD 2016)
 - ▶ curvilinear grid
 - ▶ DG, NUFFT